

Less Invasive Surfactant Administration

UC Davis Protocol

Less Invasive Surfactant Administration Protocol (LISA)

V3: 6/21/2022

Eligibility Criteria

- Preterm infants **28w0d GA to 34w0d GA**
- Infants on CPAP with FiO₂ requirement >30% to maintain saturations >90% SpO₂
- Suspected RDS with signs of respiratory distress
- CXR obtained to rule out pneumothorax

Exclusion Criteria

- Suspected or known airway or pulmonary anomaly
- Infants with persistent apnea
- Concern for pneumothorax
- Meconium aspiration syndrome

Supplies

- 16g 5.25 inch angiocath with needle discarded
- T-connector
- 3mL or 6mL syringe
- Face mask and T piece or flow-inflating bag (emergency equipment)
- Suction
- Neo View Video Laryngoscope (kept in AN2 office) or direct laryngoscope

- Paper measuring tape or use gauze package.
- Tape for marking angiocath (ETT table or cloth tape)

Surfactant

- Infasurf® (calfactant) at 3ml/kg plus 0.5ml to prime t-connector and angiocath.
- Interval between dosing: 12 hours (if needed)

Providers for procedure

- Laryngoscopy: NICU fellow, NNP, Transport RN, or NICU attending who has been trained in LISA administration
- Instillation of surfactant: Respiratory therapist, fellow/attending/NNP/transport RN
- Additional assistance: Bedside RN

Preparation for LISA

1. Prioritize non-pharmacologic approaches to analgesia such as swaddling, and sucrose solution. Use atropine to blunt reflex bradycardia. Pharmacologic sedation to be used at the discretion of the attending physician.
2. Position infant supine with CPAP/NIPPV in place, similar to positioning for standard intubation. Rotate CPAP tubing over infant's right ear to allow more space for laryngoscope insertion.
3. Ensure needle is removed from angiocath and discarded.
4. Wrap the tape around the angiocath at target depth (7cm)
 - a. Tape should indicate distance at the lip
5. If desired, bend angio-cath into hockey-stick shape to facilitate placement.
6. Draw up the surfactant (3ml/kg of calfactant + 0.5ml for priming the line and angiocath)
7. Prime the T-connector and angiocath with 0.5ml of surfactant

8. Prepare team for procedure: Read LISA Timeout Script

Performing LISA

1. With CPAP in place, perform video laryngoscopy and tracheal cannulation using angiocath.
 - a. If the CPAP is removed during visualization of airway, replace as soon as angiocath is inserted, ideally within 10 seconds.
2. Once catheter is just through the vocal cords, hold angiocath securely in position
3. Remove laryngoscope blade.
4. Connect primed T-connector/syringe to angiocath hub
5. Instill surfactant two aliquots infusions given over 10 to 30 seconds
 - a. Aspirate stomach contents after each aliquot to check for inadvertent esophageal administration of surfactant.
 - b. Anticipate small amounts of surfactant bubbling from lips with correctly placed angiocath.
6. Infant remains supine and midline, no turning necessary
7. Once instillation complete, remove angiocath and continue CPAP/NIPPV

Procedure Notes

- Please document as a procedure note in EPIC using the dot phrase .NICULISA
- If first attempt at tracheal catheterization is not possible within 60 seconds or with a desat >20 points below baseline, remove laryngoscope and allow infant to recover on CPAP.
- Total of 3 attempts allowed for LISA until need to transition to intubation for surfactant


- Increase FiO₂ 10-20% at start of procedure.
- Wait longer than 30 seconds for next bolus of surfactant if bradycardia <80bpm for > 5 seconds OR desaturation to <70% SpO₂ for >10 seconds.
- If mild apnea, can give breaths on vent, if prolonged or severe transition to PPV per NRP. Ok to give PPV while angiocath is in place. Insure mouth is closed.

Medical Legal Disclaimer:

Welcome to the UC Davis Health, Department of Pediatrics, Clinical Practice Guidelines Website. All health and health-related information contained within the Site is intended chiefly for use as a resource by the Department's clinical staff and trainees in the course and scope of their approved functions/activities (although it may be accessible by others via the internet). This Site is not intended to be used as a substitute for the exercise of independent professional judgment. These clinical pathways are intended to be a guide for practitioners and may need to be adapted for each specific patient based on the practitioner's professional judgment, consideration of any unique circumstances, the needs of each patient and their family, and/or the availability of various resources at the health care institution where the patient is located. Efforts are made to ensure that the material within this Site is accurate and timely but is provided without warranty for quality or accuracy. The Regents of the University of California; University of California, Davis; University of California, Davis, Health nor any other contributing author is responsible for any errors or omissions in any information provided or the results obtained from the use of such information. Some pages within this Site, for the convenience of users, are linked to or may refer to websites not managed by UC Davis Health. UC Davis Health does not control or take responsibility for the content of these websites, and the views and opinions of the documents in this Site do not imply endorsement or credibility of the service, information or product offered through the linked sites by UC Davis Health. UC Davis Health provides limited personal permission to use the Site. This Site is limited in that you may not:

- Use, download or print material from this site for commercial use such as selling, creating course packets, or posting information on another website.
- Change or delete propriety notices from material downloaded or printed from it. · Post or transmit any unlawful, threatening, libelous, defamatory, obscene, scandalous, inflammatory, pornographic, or profane material, any propriety information belonging to others or any material that could be deemed as or encourage criminal activity, give rise to civil liability, or otherwise violate the law.
- Use the Site in a manner contrary to any applicable law.

You should assume that everything you see or read on this Site is copyrighted by University of California or others unless otherwise noted. You may download information from this Site as long as



it is not used for commercial purposes, and you retain the proprietary notices. You may not use, modify, make multiple copies, or distribute or transmit the contents of this Site for public or commercial purposes without the express consent of UC Davis Health.