




Foreign Body Ingestion Algorithms: Button Battery, Coin/Blunt Objects, Magnet, Sharp Objects, Esophageal Food Impaction

BUTTON BATTERY INGESTION

1. Obtain Button Battery (BB) information, if possible.
2. Call National Battery Ingestion Hotline: **800-498-8666**
3. Obtain STAT X-rays: lateral neck (including nasal cavities and nasopharynx) and AP or PA chest, and AP abdomen/pelvis views.
4. Keep patient NPO except for sucralfate (as directed below). Start IV and MIVF.
5. Submit report to Consumer Product Safety Commission at www.SaferProducts.gov. See QR code below.



*Sucralfate (1g/10mL suspension): 10mL q10min, max 3 doses



Scan this QR code to make a report to the US Consumer Product Safety Commission

UC DAVIS
HEALTH

CHILDREN'S
HOSPITAL



Verified Level I
Children's Surgery
Center

Endoscopic Retrieval Tips

- Removal should be performed under general anesthesia
- Inspect mucosa for extent, location, and depth of damage

Discharge Considerations


- Bowel regimen
- Reliable/competent caregiver
- Regular diet
- Encourage activity
- Educate on signs of GI bleed (bloodstained vomiting, melena, abdominal pain, change in eating pattern) or features of intestinal obstruction (persistent vomiting, abdominal distention)

References

- [Management of Ingested Foreign Bodies in Children: A Clinical Report of the NASPGHAN Endoscopy Committee](#) Kramer, Robert E.; Lerner, Diana G.; Lin, Tom; Manfredi, Michael; Shah, Manoj; Stephen, Thomas C.; Gibbons, Troy E.; Pall, Harpreet; Sahn, Ben; McOmber, Mark; Zacur, George; Friedlander, Joel; Quiros, Antonio J.; Fishman, Douglas S.; Mamula, Petar
- Journal of Pediatric Gastroenterology and Nutrition 60(4):562-574, April 2015. doi: 10.1097/MPG.0000000000000729
- National Capital Poison Control: www.poison.org/battery/guideline
- <https://www.chop.edu/clinical-pathway/foreign-body-ingestion-clinical-pathway>

COIN & OTHER BLUNT OBJECT INGESTION

1. **Obtain history: Ensure no concern for button battery or magnet ingestion (refer to those algorithms if there is concern).**
2. Obtain STAT X-rays: lateral neck (including nasal cavities and nasopharynx) and AP or PA chest, and AP abdomen/pelvis views.
3. Keep NPO. If symptomatic or esophageal FB, start IV and MIVF.
4. Consult: ENT for above the clavicles; Peds GI for esophagus, below clavicles, or as needed for stomach or intestinal FB.
5. Submit report (if applicable) to Consumer Product Safety Commission at www.SaferProducts.gov. See QR code below.



Any concern for perforation or obstruction or history of esophageal or abdominal surgery:
Consult Pediatric Surgery

ESOPHAGUS

Symptomatic:
Drooling,
dysphagia,
respiratory
compromise,
severe pain

Timely
endoscopic
removal
depending on
symptom
severity

Asymptomatic

1. Repeat X-ray to ensure coin is still present
2. Urgent endoscopic removal

STOMACH

Symptomatic

Asymptomatic

No endoscopy needed.
Consider straining stools,
laxatives.
Repeat X-ray at 2 weeks to
confirm passage, or sooner
if symptoms develop.
Depending on size of
object, consider admission
for observation (larger
objects could get caught at
ileocecal valve).

Endoscopic removal if
not passed in 2-4
weeks.
Repeat X-ray before
removing to ensure
coin is still present.

SMALL BOWEL

Symptomatic

Consider
alternative
etiologies and
alternative
foreign bodies.
Surgical removal
if indicated.

Coin Sizes:

Quarter:
24mm
Nickel: 21mm
Penny: 19mm
Dime: 18mm



Scan this QR code to make a report to the US
Consumer Product Safety Commission

See reverse for
additional information



UC DAVIS
HEALTH

CHILDREN'S
HOSPITAL



Verified Level I
Children's Surgery
Center

Endoscopic Retrieval Tips

- Removal should be performed under general anesthesia
- Inspect mucosa for extent, location, and depth of damage

Discharge Considerations


- Bowel regimen
- Reliable/competent caregiver
- Regular diet
- Encourage activity
- Educate on signs of GI bleed (bloodstained vomiting, melena, abdominal pain, change in eating pattern) or features of intestinal obstruction (persistent vomiting, abdominal distention)

References

- [Management of Ingested Foreign Bodies in Children: A Clinical Report of the NASPGHAN Endoscopy Committee](#) Kramer, Robert E.; Lerner, Diana G.; Lin, Tom; Manfredi, Michael; Shah, Manoj; Stephen, Thomas C.; Gibbons, Troy E.; Pall, Harpreet; Sahn, Ben; McOmber, Mark; Zacur, George; Friedlander, Joel; Quiros, Antonio J.; Fishman, Douglas S.; Mamula, Petar
- Journal of Pediatric Gastroenterology and Nutrition 60(4):562-574, April 2015. doi: 10.1097/MPG.0000000000000729
- National Capital Poison Control: www.poison.org/battery/guideline
- <https://www.chop.edu/clinical-pathway/foreign-body-ingestion-clinical-pathway>

MAGNET INGESTION

1. Obtain history: known magnet ingestion?
2. Obtain STAT X-rays: lateral neck (including nasal cavities and nasopharynx) and AP or PA chest, and AP abdomen/pelvis views : Determine single versus multiple magnets.
3. Keep NPO. Start IV and MIVF.
4. Consult: ENT for above the clavicles; Peds GI for esophagus, stomach, intestine, or below clavicles.
5. Submit report to Consumer Product Safety Commission at www.SaferProducts.gov. See QR code below.



Scan this QR code to make a report to the US Consumer Product Safety Commission

UC DAVIS
HEALTH

CHILDREN'S
HOSPITAL



Verified Level I
Children's Surgery
Center

See reverse for additional information

Endoscopic Retrieval Tips

- Removal should be performed under general anesthesia
- Inspect mucosa for extent, location, and depth of damage

Discharge Considerations- When Waiting for Single Magnet to Pass


- Bowel regimen
- Reliable/competent caregiver
- Regular diet
- Encourage activity
- Educate on signs of GI bleed (bloodstained vomiting, melena, abdominal pain, change in eating pattern) or features of intestinal obstruction (persistent vomiting, abdominal distention)
- Remove any magnetic objects nearby
- Avoid clothes with metallic buttons or belts with buckles
- Ensure no other metal objects or magnets are in the child's environment for accidental ingestion
- Discuss notification of Product Safety Commission

References

- [Management of Ingested Foreign Bodies in Children: A Clinical Report of the NASPGHAN Endoscopy Committee](#) Kramer, Robert E.; Lerner, Diana G.; Lin, Tom; Manfredi, Michael; Shah, Manoj; Stephen, Thomas C.; Gibbons, Troy E.; Pall, Harpreet; Sahn, Ben; McOmber, Mark; Zacur, George; Friedlander, Joel; Quiros, Antonio J.; Fishman, Douglas S.; Mamula, Petar
- Journal of Pediatric Gastroenterology and Nutrition 60(4):562-574, April 2015. doi: 10.1097/MPG.0000000000000729
- National Capital Poison Control: www.poison.org/battery/guideline
- <https://www.chop.edu/clinical-pathway/foreign-body-ingestion-clinical-pathway>

SHARP OBJECT INGESTION

1. Obtain STAT X-rays: lateral neck (including nasal cavities and nasopharynx) and AP or PA chest, and AP abdomen/pelvis views.
2. Keep NPO. If esophageal or symptomatic, start IV and MIVF.
3. Consult: ENT for above the clavicles; Peds GI for esophagus, intestine, or below clavicles.
4. Submit report (if applicable) to Consumer Product Safety Commission at www.SaferProducts.gov. See QR code below.



See reverse for additional information →



Scan this QR code to make a report to the US Consumer Product Safety Commission

UC DAVIS
HEALTH

CHILDREN'S
HOSPITAL



Verified Level I
Children's Surgery
Center

Endoscopic Retrieval Tips

- Removal should be performed under general anesthesia
- Inspect mucosa for extent, location, and depth of damage
- If sharp end is facing cephalad, consider pushing the object into the stomach and rotate the sharp end caudally before removal
- Safety pin: polypectomy snare can be used to close it before withdrawal
- Toothpick: polypectomy snare is a good option

Discharge Considerations


- Reliable/competent caregiver
- Regular diet
- Encourage activity
- Educate on signs of GI bleed (bloodstained vomiting, melena, abdominal pain, change in eating pattern) or features of intestinal obstruction (persistent vomiting, abdominal distention)

References

- [Management of Ingested Foreign Bodies in Children: A Clinical Report of the NASPGHAN Endoscopy Committee](#) Kramer, Robert E.; Lerner, Diana G.; Lin, Tom; Manfredi, Michael; Shah, Manoj; Stephen, Thomas C.; Gibbons, Troy E.; Pall, Harpreet; Sahn, Ben; McOmber, Mark; Zacur, George; Friedlander, Joel; Quiros, Antonio J.; Fishman, Douglas S.; Mamula, Petar
- Journal of Pediatric Gastroenterology and Nutrition 60(4):562574, April 2015. doi: 10.1097/MPG.0000000000000729
- National Capital Poison Control: www.poison.org/battery/guideline
- <https://www.chop.edu/clinical-pathway/foreign-body-ingestion-clinical-pathway>

ESOPHAGEAL FOOD IMPACTION

1. Obtain STAT X-rays: lateral neck (including nasal cavities and nasopharynx) and AP or PA chest, and AP abdomen/pelvis views.
2. Keep NPO. Start IV and MIVF.
3. Consult Peds GI.



See reverse for
additional information



**UC DAVIS
HEALTH**

**CHILDREN'S
HOSPITAL**



Verified Level I
Children's Surgery
Center

Endoscopic Retrieval Tips

- Removal should be performed under general anesthesia
- Inspect mucosa for extent, location, and depth of damage
- Obtain proximal and distal esophageal biopsies
- Assess for stricture. Dilation is cautioned against if eosinophilic esophagitis is suspected, or food has been impacted for a long duration

Discharge Considerations

- Ensure patient has appropriate GI followup

References

- [Management of Ingested Foreign Bodies in Children: A Clinical Report of the NASPGHAN Endoscopy Committee](#) Kramer, Robert E.; Lerner, Diana G.; Lin, Tom; Manfredi, Michael; Shah, Manoj; Stephen, Thomas C.; Gibbons, Troy E.; Pall, Harpreet; Sahn, Ben; McOmber, Mark; Zacur, George; Friedlander, Joel; Quiros, Antonio J.; Fishman, Douglas S.; Mamula, Petar
- Journal of Pediatric Gastroenterology and Nutrition 60(4):562574, April 2015. doi: 10.1097/MPG.0000000000000729
- National Capital Poison Control: www.poison.org/battery/guideline
- <https://www.chop.edu/clinical-pathway/foreign-body-ingestion-clinical-pathway>

Medical Legal Disclaimer:

Welcome to the UC Davis Health, Department of Pediatrics, Clinical Practice Guidelines Website. All health and health-related information contained within the Site is intended chiefly for use as a resource by the Department's clinical staff and trainees in the course and scope of their approved functions/activities (although it may be accessible by others via the internet). This Site is not intended to be used as a substitute for the exercise of independent professional judgment. These clinical pathways are intended to be a guide for practitioners and may need to be adapted for each specific patient based on the practitioner's professional judgment, consideration of any unique circumstances, the needs of each patient and their family, and/or the availability of various resources at the health care institution where the patient is located. Efforts are made to ensure that the material within this Site is accurate and timely but is provided without warranty for quality or accuracy. The Regents of the University of California; University of California, Davis; University of California, Davis, Health nor any other contributing author is responsible for any errors or omissions in any information provided or the results obtained from the use of such information. Some pages within this Site, for the convenience of users, are linked to or may refer to websites not managed by UC Davis Health. UC Davis Health does not control or take responsibility for the content of these websites, and the views and opinions of the documents in this Site do not imply endorsement or credibility of the service, information or product offered through the linked sites by UC Davis Health. UC Davis Health provides limited personal permission to use the Site. This Site is limited in that you may not:

- Use, download or print material from this site for commercial use such as selling, creating course packets, or posting information on another website.
- Change or delete propriety notices from material downloaded or printed from it. · Post or transmit any unlawful, threatening, libelous, defamatory, obscene, scandalous, inflammatory, pornographic, or profane material, any propriety information belonging to others or any material that could be deemed as or encourage criminal activity, give rise to civil liability, or otherwise violate the law.
- Use the Site in a manner contrary to any applicable law.

You should assume that everything you see or read on this Site is copyrighted by University of California or others unless otherwise noted. You may download information from this Site as long as it is not used for commercial purposes, and you retain the proprietary notices. You may not use, modify, make multiple copies, or distribute or transmit the contents of this Site for public or commercial purposes without the express consent of UC Davis Health.