

Pediatric Home NGT Guidelines

Background:

There are many reasons why an older infant (>60weeks CGA) or child may not be able to achieve full oral feeds or may lose the ability to feed orally for a period of time. For some children requiring short-term enteral nutrition support (typically <3 months), a nasogastric tube may be the most appropriate route for nutritional support. For other children that will likely need enteral feeding support for longer durations (typically >3 months), a G-Tube may be the appropriate choice.

Criteria for Home NGT:

- Patient has demonstrated appropriate growth on anticipated home regiment for >48 hours
- Caregiver(s) are agreeable to home NGT and have completed all training requirements
- Caregiver(s) have access to outpatient follow up with PCP comfortable managing NGT and/or a primary pediatric subspecialty team with RD support with scheduled follow-up
- Caregiver(s) have access to home equipment resources (DME).
- Caregivers are able to demonstrate all cares required for home NGT feeds and medication administration. This may include 24-48 hours of rooming in stay.

Caregiver Training:

- 2 available caregivers identified
- Each caregiver must demonstrate 1-2 placements on child, and additional training can be completed on a manikin. This will be also determined by age and/or nutritional needs of patient (ie. risk of delayed feeds, hypoglycemia, etc.)
- Caregiver can connect NGT to feeding source (Gravity vs Pump).
- If patients go home with Feeding Pump, able to demonstrate how to operate the pump.
- Determine safety risks for consideration of overnight or prolonged feeds via a pump.

Discharge Medical Equipment Needed:

- Use Order Set: "Neonatal/Pediatric Feeding DME". This includes NG/NJ, G tube, and GJ discharge DME needs.

NG securement devices:

- At times, a NG securement device called a bridle may be the most appropriate way to ensure displacement of the NG. This may be helpful with a critical NG tube such as in the setting of esophageal stricture or caustic ingestion where the NG tube is ensuring patency of the esophagus as well as a route for feeding.

Outpatient Follow Up:

- Communicate with PCP prior to discharge. Ensure PCP is comfortable with initial management
- Discuss routine replacement and dislodgement plan with caregivers.
- Identify which outpatient dietitian and subspecialty team will be managing feeds and ensure active referral in place and communicate timing of discharge to ensure follow-up
-

References:

1. Singhal S, Baker SS, Bojczuk GA, Baker RD. Tube Feeding in Children. *Pediatr Rev.* 2017 Jan;38(1):23-34.
2. Abdelhadi RA, Rempel G, Sevilla W, Turner JM, Quet J, Nelson A, Rahe K, Wilhelm R, Larocque J, Guenter P; ASPEN Enteral Nutrition Task Force Pediatric Work Group. Transitioning From Nasogastric Feeding Tube to Gastrostomy Tube in Pediatric Patients: A Survey on Decision-Making and Practice. *Nutr Clin Pract.* 2021 Jun;36(3):654-664.
3. Khalil ST, Uhing MR, Duesing L, Visotcky A, Tarima S, Nghiem-Rao TH. Outcomes of Infants With Home Tube Feeding: Comparing Nasogastric vs Gastrostomy Tubes. *JPEN J Parenter Enteral Nutr.* 2017 Nov;41(8):1380-1385.

Medical Legal Disclaimer:

Welcome to the UC Davis Health, Department of Pediatrics, Clinical Practice Guidelines Website. All health and health-related information contained within the Site is intended chiefly for use as a resource by the Department's clinical staff and trainees in the course and scope of their approved functions/activities (although it may be accessible by others via the internet). This Site is not intended to be used as a substitute for the exercise of independent professional judgment. These clinical pathways are intended to be a guide for practitioners and may need to be adapted for each specific patient based on the practitioner's professional judgment, consideration of any unique circumstances, the needs of each patient and their family, and/or the availability of various resources at the health care institution where the patient is located. Efforts are made to ensure that the material within this Site is accurate and timely but is provided without warranty for quality or accuracy. The Regents of the University of California; University of California, Davis; University of California, Davis, Health nor any other contributing author is responsible for any errors or omissions in any information provided or the results obtained from the use of such information. Some pages within this Site, for the convenience of users, are linked to or may refer to websites not managed by UC Davis Health. UC Davis Health does not control or take responsibility for the content of these websites, and the views and opinions of the documents in this Site do not imply endorsement or credibility of the service, information or product offered through the linked sites by UC Davis Health. UC Davis Health provides limited personal permission to use the Site. This Site is limited in that you may not:

- Use, download or print material from this site for commercial use such as selling, creating course packets, or posting information on another website.
- Change or delete propriety notices from material downloaded or printed from it. · Post or transmit any unlawful, threatening, libelous, defamatory, obscene, scandalous, inflammatory, pornographic, or profane material, any propriety information belonging to others or any material that could be deemed as or encourage criminal activity, give rise to civil liability, or otherwise violate the law.
- Use the Site in a manner contrary to any applicable law.

You should assume that everything you see or read on this Site is copyrighted by University of California or others unless otherwise noted. You may download information from this Site as long as it is not used for commercial purposes, and you retain the proprietary notices. You may not use, modify, make multiple copies, or distribute or transmit the contents of this Site for public or commercial purposes without the express consent of UC Davis Health.